



**UNITED STATES
CONSUMER PRODUCT SAFETY COMMISSION
4330 EAST WEST HIGHWAY
BETHESDA, MD 20814**

Memorandum

This document has been electronically
approved and signed.

Date: January 25, 2021

TO : Duane Boniface, Director
Office of Hazard Identification and Reduction

THROUGH : Risana Chowdhury, Director
Division of Hazard Analysis

FROM : Angie Qin, Mathematical Statistician
Division of Hazard Analysis

Cheryl Scorpio, Ph.D., Pharmacologist
Directorate for Health Sciences

SUBJECT : Unintentional Pediatric Poisoning Injury Estimates for 2019

In 2019, there were an estimated 67,500 emergency department (ED)-treated injuries involving unintentional pediatric poisonings.¹ Unintentional pediatric poisonings are poisonings and chemical burns² resulting from accidental access to a chemical substance by a child under the age of 5 years old. Adverse reactions, therapeutic errors, and incidents that are not addressable through the Poison Prevention Packaging Act (PPPA) were not included in the estimates.

Results

Staff found 2,450 cases involving unintentional pediatric poisonings in 2019 in the National Electronic Injury Surveillance System (NEISS). Based on these cases, staff computed a national estimate of 67,500 ED-treated injuries, with a coefficient of variance (C.V.) of 13.24 percent. The 95 percent confidence interval (C.I.) for this estimate was 50,000 to 85,000. A breakdown of the estimates by diagnosis is shown in Table 1.

¹ In October 2018, an upgrade was implemented for NEISS. An ED visit is allowed to contain up to two codes for the diagnoses. Data were extracted if either of the two codes listed poisoning or chemical burn. Note the diagnosis categories are not mutually exclusive because it is possible for some incidents to have both poisoning and chemical burn diagnoses.

² Chemical burns are included in this memorandum because many of the substances regulated by the PPPA cause chemical burns. Examples of such substances include: tire cleaners, etching creams, drain cleaners, and oven cleaners.

*This analysis was prepared by CPSC staff and has not been reviewed or approved by,
and may not necessarily reflect the views of, the Commission.*

CPSC Hotline: 1-800-638-CPSC (2772) - CPSC's Web Site: <http://www.cpsc.gov>

THIS DOCUMENT HAS NOT BEEN REVIEWED
OR ACCEPTED BY THE COMMISSION

CLEARED FOR PUBLIC RELEASE
UNDER CPSA 6(b)(1)

Table 1: 2019 Emergency Department-Treated Unintentional Pediatric Poisoning Estimates by Diagnosis*

Diagnosis (Code)	Estimate	Cases	C.V.	95% C.I.
Poisoning (68)	64,500	2,356	12.21%	49,100-80,000
Chemical Burn (49)	N/A**	100	41.21%	N/A**
Total ³	67,500	2,450	13.24%	50,000-85,000

Source: National Electronic Injury Surveillance System, June 2020

* Adjusted to exclude adverse reactions, therapeutic errors, and exposures beyond the victim's control.⁴

** Estimates are unstable as c.v. is above the threshold of 33%.

Table 2 gives a breakdown by year of the estimated ED-treated unintentional pediatric poisonings. The estimates decreased from 2017 to 2019, but the trend was not statistically significant (the lowest p-value for all trends was 0.25).

Table 2: 2017–2019 Emergency Department Treated Unintentional Pediatric Poisoning Estimates by Year*

Diagnosis (Code)	2017	2018	2019	Average
Poisoning (68)	79,000	66,800	64,500	70,100
Chemical Burn (49)	4,100	4,200	N/A**	N/A**
Total ³	83,200	70,900	67,500	73,800

Source: National Electronic Injury Surveillance System, June 2020

* Adjusted to exclude adverse reactions, therapeutic errors, and exposures beyond the victim's control.

** Estimates are unstable as c.v. is above the threshold of 33%.

In 2019, an estimated 57,400 (85 percent of the total 67,500) ED-treated unintentional pediatric poisonings occurred at home. An estimated 9,100 (13 percent) poisonings occurred at an unknown location. The remaining injuries occurred at other locations, such as streets, schools, playgrounds, and other public property.

Table 3 shows the rolling 3-year average of the estimated ED-treated unintentional pediatric poisonings from 2013 to 2019. The somewhat higher average estimate for chemical burns during 2013–2015 was mainly due to laundry packet chemical burn incidents. The change was not statistically significant.

Table 3: Rolling 3-Year Average for Emergency Department-Treated Unintentional Pediatric Poisoning Estimates from 2013 to 2019*

Diagnosis (Code)	Average (2013-2015)	Average (2014-2016)	Average (2015-2017)	Average (2016-2018)	Average (2017-2019)
Poisoning (68)	80,700	80,800	79,400	74,200	70,100
Chemical Burn (49)	3,900	3,700	3,600	3,700	N/A**
Total ³	84,600	84,500	83,000	77,900	73,800

Source: National Electronic Injury Surveillance System, June 2020

* Adjusted to exclude adverse reactions, therapeutic errors, and exposures beyond the victim's control.

** Estimates are unstable as c.v. is above the threshold of 33%.

³ Columns may not sum to totals due to rounding and the diagnosis categories are not mutually exclusive

⁴ See examples of the out-of-scope cases in Methodology section on page 4.

Table 4 gives a breakdown, by the product involved, for the estimated ED-treated unintentional pediatric poisonings. Note that the product categories are not mutually exclusive because it is possible for two different products to be associated with the same poisoning incident.

Table 4: 2019 Emergency Department Treated Unintentional Pediatric Poisoning Estimates by Top 10 Products^{5*}

Product	Estimate ⁶	C.V.	95% C.I.
Blood Pressure Medications	7,400	17.08%	4,900-9,900
Acetaminophen	5,700	13.56%	4,200-7,200
Bleach	2,800	28.33%	1,300-4,400
Ibuprofen	2,700	21.50%	1,600-3,900
Antidepressants	2,600	17.28%	1,700-3,500
Attention Deficit Disorder Medications	2,600	23.40%	1,400-3,800
Laundry Packets	2,500	20.88%	1,500-3,600
Dietary Supplements	2,400	20.51%	1,500-3,400
Diphenhydramine	2,300	21.42%	1,400-3,300
Unknown	3,700	18.74%	2,300-5,000

Source: National Electronic Injury Surveillance System, June 2020

** Adjusted to exclude adverse reactions, therapeutic errors, and exposures beyond the victim's control*

Table 5 shows that the ED-treated unintentional pediatric poisoning estimates decreased for Acetaminophen, bleach, Ibuprofen and laundry packets, increased for blood pressure medications, antidepressants and unknown medications, and remained unchanged for Diphenhydramine, from calendar year 2018 to 2019. Attention deficit disorder medications and dietary supplements moved into the top 10 in 2019. Narcotics medications and household cleaners dropped out of the top 10 in 2019.

Table 5: 2018–2019 Emergency Department-Treated Unintentional Pediatric Poisoning Estimates by Top 10 Products*

2018	2019
Blood Pressure Medications (6,300)	Blood Pressure Medications (7,400)
Acetaminophen (6,000)	Acetaminophen (5,700)
Bleach (3,900)	Bleach (2,800)
Ibuprofen (3,300)	Ibuprofen (2,700)
Laundry Packets (2,700)	Antidepressants (2,600)
Diphenhydramine (2,300)	Attention Deficit Disorder Meds (2,600)
Antidepressants (2,300)	Laundry Packets (2,500)
Narcotics Medications (2,000)	Dietary Supplements (2,400)
Household Cleaners (1,900)	Diphenhydramine (2,300)
Unknown (2,700)	Unknown (3,700)

Source: National Electronic Injury Surveillance System, June 2020

** Adjusted to exclude adverse reactions, therapeutic errors, and exposures beyond the victim's control*

⁵ Liquid nicotine poisoning injury did not appear in the top 10 products in 2019, or any prior years. The Child Nicotine Poisoning Prevention Act of 2015 (CNPPA), requires any nicotine provided in a liquid nicotine container, sold, offered for sale, manufactured for sale, distributed in commerce, or imported into the United States shall be packaged in accordance with the standards of the Poison Prevention Packaging Act (PPPA).

⁶ Please refer to appendix for the estimates for prior years.

Methodology

NEISS is a probability sample of approximately 100 U.S. hospitals, each operating 24-hour EDs and providing more than six beds. Staff in each hospital input and code consumer product-related data from the ED record, and then the data are transmitted electronically to the CPSC. Because NEISS is a probability sample, each case collected represents a number of cases (the case's *weight*) in the total estimate of injuries in the United States. Different hospitals carry different weights, based on stratification by their annual number of ED visits (Kessler and Schroeder, 1999).

Hazard Analysis staff searched NEISS databases for all incidents with poisoning diagnosis (code 68) or chemical burn diagnosis (code 49) involving children under the age of 5. Health Sciences staff examined all incidents to identify cases that were not unintentional exposures, but were deemed generally associated with a prescribed therapeutic regimen, or an unforeseen incidental exposure from a situation beyond the victim's control. These types of cases, delineated below, are out-of-scope cases because they do not directly involve a child independently accessing a poison.


1. *Adverse Reactions*: This includes undesirable effects that occur with the proper use of a substance (e.g., drowsiness after administration of an antihistamine). Allergic, hypersensitivity, or idiosyncratic reactions to recommended doses of vaccines, antibiotics, or other medications are also included in this category.
2. *Therapeutic Errors*: Unintentional mistakes made during a prescribed or recommended course of treatment, such as: (1) a caregiver administering the wrong substance or an overdose (e.g., two tablespoons instead of two teaspoons) to the patient; (2) a pharmacist mislabeling the dosage instructions on a prescription; or (3) a caregiver giving medication to the wrong child.
3. *Incidental Exposures*: This category refers to exposures resulting from a situation beyond the control of the victim. Examples include exposures to: (1) chlorine fumes from a pool; (2) gas fumes while in a dwelling or an automobile; (3) gasoline while it is being pumped into an automobile; or (4) illicit drugs (e.g., cocaine, methamphetamine, marijuana) while the caregiver is using or producing them.

Hazard Analysis staff used SAS[®] software program version 9.4 to manage and retrieve data and to compute estimates and the associated C.V. for the number of unintentional pediatric poisoning injuries. A C.V. is the ratio of the standard error of the estimate (i.e., variability) to the estimate itself. This is generally expressed as a percent. A C.V. of 10 percent means the standard error of the estimate equals 0.1 times the estimate.

NEISS data do not typically identify all of the contributing factors to unintentional pediatric poisoning injuries. CPSC continues public outreach efforts to help manufacturers comply with the PPPA and to remind consumers about the need to keep products in their original child-resistant packaging and out of the reach of children.

Appendix A

Figure A.1: Estimates for Products in the Top 10 Every Year from 2011 to 2019*




**Due to limited variance in the number of children under 5, the pattern remains the same when normalizing.*

**There is a reduction in Acetaminophen since 2011, which may be due to several Acetaminophen child safety cap recalls*

Source: National Electronic Injury Surveillance System

**Figure A.2: Estimates for Other Products (Laundry Packet and Antiepileptics)
From 2011 to 2019***



** Laundry Packet was introduced in United States in 2012*

Source: National Electronic Injury Surveillance System

Table A.1: 2011 Emergency Department Treated Unintentional Pediatric Poisoning Estimates by Top 10 Products*

Product	Estimate	C.V.	95% C.I.
Acetaminophen	8,400	13.81%	6,100-10,700
Blood Pressure Medications	6,600	14.81%	4,700-8,600
Bleach	4,400	15.20%	3,100-5,700
Ibuprofen	3,800	14.37%	2,700-4,800
Sedatives and Antianxiety Medications	3,600	18.61%	2,300-4,900
Vitamins	3,200	17.07%	2,100-4,300
Antidepressants	2,900	15.86%	2,000-3,900
Muscle Relaxants	2,800	19.14%	1,800-3,900
Narcotics Medications	2,500	22.63%	1,400-3,700
Unknown	4,600	15.73%	3,200-6,000

Source: National Electronic Injury Surveillance System

** Adjusted to exclude adverse reactions, therapeutic errors, and exposures beyond the victim's control*

Table A.2: 2012 Emergency Department Treated Unintentional Pediatric Poisoning Estimates by Top 10 Products*

Product	Estimate	C.V.	95% C.I.
Acetaminophen	9,400	14.05%	6,800-12,000
Blood Pressure Medications	6,200	15.13%	4,400-8,100
Sedatives and Antianxiety Medications	4,100	17.86%	2,600-5,500
Antidepressants	3,700	20.18%	2,200-5,100
Narcotics Medications	3,700	15.80%	2,500-4,800
Bleach	3,600	18.82%	2,300-5,000
Ibuprofen	3,400	14.65%	2,400-4,400
Diphenhydramine	2,700	14.19%	2,000-3,500
Household Cleaners	2,700	14.26%	1,900-3,400
Unknown	4,100	14.84%	2,900-5,300

Source: National Electronic Injury Surveillance System

** Adjusted to exclude adverse reactions, therapeutic errors, and exposures beyond the victim's control*

Table A.3: 2013 Emergency Department-Treated Unintentional Pediatric Poisoning Estimates by Top 10 Products*

Product	Estimate	C.V.	95% C.I.
Acetaminophen	8,100	14.50%	5,800-10,400
Blood Pressure Medications	6,500	15.55%	4,500-8,500
Bleach	3,100	15.12%	2,200-4,100
Antidepressants	3,100	21.38%	1,800-4,400
Ibuprofen	3,000	17.42%	2,000-4,100
Laundry Packets	3,000	17.91%	1,900-4,000
Sedatives and Antianxiety Medications	2,700	27.82%	1,200-4,100
Narcotics Medications	2,400	19.68%	1,500-3,300
Diphenhydramine	2,300	19.93%	1,400-3,100
Unknown	4,100	18.64%	2,600-5,500

Source: National Electronic Injury Surveillance System

** Adjusted to exclude adverse reactions, therapeutic errors, and exposures beyond the victim's control*

Table A.4: 2014 Emergency Department Treated Unintentional Pediatric Poisoning Estimates by Top 10 Products*

Product	Estimate	C.V.	95% C.I.
Blood Pressure Medications	7,300	17.67%	4,800-9,800
Acetaminophen	6,800	14.24%	4,900-8,600
Bleach	4,700	15.88%	3,300-6,200
Laundry Packets	4,100	20.96%	2,400-5,800
Antidepressants	3,400	16.01%	2,300-4,500
Diphenhydramine	3,200	20.50%	1,900-4,500
Narcotics Medications	3,200	25.45%	1,600-4,800
Opioid Antagonists	3,100	28.29%	1,400-4,800
Ibuprofen	2,900	19.61%	1,800-4,000
Unknown	3,500	18.05%	2,300-4,800

Source: National Electronic Injury Surveillance System

** Adjusted to exclude adverse reactions, therapeutic errors, and exposures beyond the victim's control*

Table A.5: 2015 Emergency Department Treated Unintentional Pediatric Poisoning Estimates by Top 10 Products*

Product	Estimate	C.V.	95% C.I.
Acetaminophen	7,900	14.49%	5,700-10,200
Blood Pressure Medications	7,300	16.22%	5,000-9,700
Laundry Packets	5,000	18.63%	3,100-6,800
Ibuprofen	3,700	16.82%	2,500-4,900
Bleach	3,400	19.04%	2,100-4,600
Antidepressants	2,800	18.14%	1,800-3,800
Narcotics Medications	2,700	21.01%	1,600-3,800
Sedatives and Antianxiety Medications	2,600	17.76%	1,700-3,600
Diphenhydramine	2,500	18.67%	1,600-3,400
Unknown	3,100	15.23%	2,200-4,000

Source: National Electronic Injury Surveillance System

** Adjusted to exclude adverse reactions, therapeutic errors, and exposures beyond the victim's control*

Table A.6: 2016 Emergency Department Treated Unintentional Pediatric Poisoning Estimates by Top 10 Products*

Product	Estimate	C.V.	95% C.I.
Blood Pressure Medications	6,000	19.64%	3,700-8,300
Acetaminophen	5,900	16.21%	4,000-7,700
Laundry Packets	3,600	19.18%	2,200-4,900
Bleach	3,000	19.83%	1,800-4,200
Sedatives and Antianxiety Medications	2,800	19.70%	1,700-3,900
Antidepressants	2,600	21.15%	1,500-3,600
Ibuprofen	2,500	24.52%	1,300-3,800
Narcotic Medications	2,400	22.93%	1,300-3,400
Diphenhydramine	2,400	21.32%	1,400-3,300
Unknown	3,600	18.85%	2,300-5,000

Source: National Electronic Injury Surveillance System

** Adjusted to exclude adverse reactions, therapeutic errors, and exposures beyond the victim's control*

Table A.7: 2017 Emergency Department Treated Unintentional Pediatric Poisoning Estimates by Top 10 Products*

Product	Estimate	C.V.	95% C.I.
Blood Pressure Medications	6,700	21.22%	3,900-9,400
Acetaminophen	6,400	14.21%	4,600-8,200
Bleach	3,500	21.98%	2,000-5,100
Ibuprofen	3,400	20.67%	2,000-4,700
Laundry Packets	3,300	21.62%	1,900-4,800
Antidepressants	3,100	25.80%	1,500-4,700
Vitamins	2,900	18.77%	1,900-4,000
Diphenhydramine	2,700	22.70%	1,500-3,900
Antiepileptics	2,400	26.21%	1,200-3,700
Unknown	3,500	17.64%	2,300-4,700

Source: National Electronic Injury Surveillance System

** Adjusted to exclude adverse reactions, therapeutic errors, and exposures beyond the victim's control*

Table A.8: 2018 Emergency Department Treated Unintentional Pediatric Poisoning Estimates by Top 10 Products*

Product	Estimate	C.V.	95% C.I.
Blood Pressure Medications	6,300	16.69%	4,300-8,400
Acetaminophen	6,000	15.08%	4,200-7,700
Bleach	3,900	19.80%	2,400-5,400
Ibuprofen	3,300	17.78%	2,100-4,400
Laundry Packets	2,700	17.41%	1,800-3,600
Diphenhydramine	2,300	23.97%	1,200-3,400
Antidepressants	2,300	23.29%	1,300-3,400
Household Cleaners	2,000	22.23%	1,100-2,900
Narcotics Medications	1,900	18.55%	1,200-2,600
Unknown	2,700	15.84%	1,900-3,500

Source: National Electronic Injury Surveillance System

** Adjusted to exclude adverse reactions, therapeutic errors, and exposures beyond the victim's control*